

CONVERGENCE IN CAPACITY ON COMPACT KÄHLER MANIFOLDS

SLAWOMIR DINEW, PHẠM HOÀNG HIỆP

ABSTRACT. The aim of this note is to study the convergence in capacity for functions in the class $\mathcal{E}(X, \emptyset)$. We obtain several stability theorems. Some of these are (optimal) generalizations of results of Xing [Xi2], while others are new.

INTRODUCTION

The notion of relative capacity C_n was introduced by Bedford and Taylor (cf. [BT2]). The convergence with respect to C_n in the setting of domains in $\mathbf{C}^n$ was investigated by Xing and Cegrell (see [Xi1], [Ce3]). This topic has attracted much interest, since the complex Monge-Ampère operator is continuous with respect to such a convergence (while continuity fails for, say, convergence in L^p , $1 < p < \infty$, see [CK1]). Recently Kołodziej (see [Ko1]) introduced the capacity $C_{X,\omega}$ on a compact Kähler manifold (X, ω) , which is modelled on C_n . In particular he proved that $C_{X,\omega}$ is locally equivalent to C_n . This capacity was also studied by Guedj and Zeriahi in [GZ1]. In [GZ2] they introduced the new Cegrell class $\mathcal{E}(X, \omega)$ of ω -psh functions for which the complex Monge-Ampère operator is well-defined. Roughly speaking, it is the largest class of ω -psh functions where the Monge-Ampère operator behaves like it does in the bounded functions setting.

The aim of the present note is to study the convergence in capacity $C_{X,\omega}$ in the class $\mathcal{E}(X, \omega)$ (Section 2). Thus this note might be seen as a Kähler manifold counterpart of [CK2]. Some of our results are known for subclasses of $\mathcal{E}(X, \omega)$, (especially for bounded or $\mathcal{E}^1$ functions - see [Ko2] and [Xi2]), or under additional technical assumptions (cf. [Xi2]). By applying some new results borrowed from [Di2] and [Di3] we are able to obtain the sharp results and relax those conditions.

Some of our results, however, (as well as the techniques used) are completely new, thus these might be of independent interest.

Acknowledgments. The first named author was partially supported by Polish ministerial grant N N 201 271135.

1. PRELIMINARIES First we recall some elements of pluripotential theory that will be used throughout the paper. All this can be found in [Bl], [BT1-2], [Ce1-3], [CGZ], [Di1-3], [GZ1-2], [H1-2], [Hö], [KH], [Ko1] and [Xi1-2].

1.1. Let X be a compact Kähler manifold with a fundamental form $\omega = \omega_X$ with $\int_X \omega^n = 1$. An upper semicontinuous function $\varphi : X \rightarrow [-\infty, +\infty)$ is called ω -plurisubharmonic (ω -psh for short) if $\varphi \in L^1(X)$ and $\omega_\varphi := \omega + dd^c \varphi \geq 0$, where the inequality

2000 *Mathematics Subject Classification.* 32W20, 32Q15.

Key words and phrases. Monge-Ampère operator, Cegrell classes, $\emptyset$ -plurisubharmonic functions, convergence in capacity, compact Kähler manifold.

is understood in the sense on currents. By $PSH(X, \omega)$ (resp. $PSH^-(X, \omega)$) we denote the set of ω -psh (resp. negative ω -psh) functions on X .

1.2. In [Ko1] Kołodziej introduced the capacity $C_{X, \emptyset}$ on X by

$$C_X(E) = C_{X, \emptyset}(E) = \sup \left\{ \int_E \phi_\varphi^n : \varphi \in PSH(X, \emptyset), -1 \leq \varphi \leq 0 \right\}$$

for any Borel set $E \subset X$, where $\phi_\varphi^n = (\emptyset + dd^c \varphi)^n$ and $n = \dim X$.

We refer to [Ko1], [GZ1] for more informations about this capacity.

1.3. The following class of $\emptyset$ -psh fuctions was introduced by Guedj and Zeriahi in [GZ2]:

$$\mathcal{E}(X, \emptyset) = \{ \varphi \in PSH(X, \emptyset) : \lim_{j \rightarrow \infty} \int_{\{\varphi > -j\}} \phi_{\max(\varphi, -j)}^n = \int_X \phi^n = 1 \}.$$

Intuitively it consists of those functions which have mild singularities (or do not have $-\infty$ poles at all), so that no Monge-Ampère mass concentrates near those poles. In particular any such function must have zero Lelong numbers everywhere.

Let us also define

$$\mathcal{E}^-(X, \emptyset) = \mathcal{E}(X, \emptyset) \cap PSH^-(X, \emptyset).$$

We refer to [GZ2] for all the properties of functions from $\mathcal{E}(X, \emptyset)$.

1.4. Let $u_j, u \in PSH(X, \emptyset)$. We say that $\{u_j\}$ converges to u in C_X if

$$C_X(\{|u_j - u| > \delta\}) \rightarrow 0$$

as $j \rightarrow \infty$, for every (fixed) $\delta > 0$.

1.5. A family of positive measures $\{\mu_\alpha\}$ on X is said to be uniformly absolutely continuous with respect to C_X -capacity if for every $\epsilon > 0$ there exists $\delta > 0$ such that for each Borel subset $E \subset X$ satisfying $C_X(E) < \delta$ the inequality $\mu_\alpha(E) < \epsilon$ holds for all α . We denote this by $\mu_\alpha \ll C_X$ uniformly for α . In particular all such measures must vanish on pluripolar sets.

Next we state some well-known results needed for our work. We shall sketch some of the proofs for the sake of completeness:

1.6. Proposition. *Let $u \in \mathcal{E}(X, \emptyset)$, $v \in PSH(X, \emptyset)$. Then*

$$\int_{\{u \leq v\}} \phi_{\max(u, v)}^k \wedge T = \int_{\{u \leq v\}} \phi_u^k \wedge T,$$

for all $1 \leq k \leq n$ and $T = \phi_{\varphi_1} \wedge \dots \wedge \phi_{\varphi_{n-k}}$ with $\varphi_1, \dots, \varphi_{n-k} \in \mathcal{E}(X, \emptyset)$.

Proof. By Corollary 1.7 in [GZ2] we get

$$\begin{aligned} \int_{\{u \leq v\}} \phi_{\max(u, v)}^k \wedge T &= \int_X \phi_{\max(u, v)}^k \wedge T - \int_{\{u > v\}} \phi_{\max(u, v)}^k \wedge T \\ &= \int_X \phi_u^k \wedge T - \int_{\{u > v\}} \phi_u^k \wedge T = \int_{\{u \leq v\}} \phi_u^k \wedge T. \end{aligned}$$

1.7. Proposition. *Let $u \in PSH^-(X, \emptyset)$. Then for $t \geq 0$*

$$C_X(\{u < -t\}) \leq \frac{|\sup_X u| + c}{t},$$

where the positive constant c does not depend on u .

Proof.

$$\begin{aligned}
C_X(\{u < -t\}) &= \sup_{\{u < -t\}} \left\{ \int \phi_\varphi^n : \varphi \in PSH(X, \emptyset), -1 \leq \varphi \leq 0 \right\} \\
&\leq \sup_{\{u < -t\}} \left\{ \int \frac{|u|}{t} \phi_\varphi^n : \varphi \in PSH(X, \emptyset), -1 \leq \varphi \leq 0 \right\} \\
&\leq \frac{1}{t} \sup_X \left\{ \int |u| \phi_\varphi^n : \varphi \in PSH(X, \emptyset), -1 \leq \varphi \leq 0 \right\} \\
&\leq \frac{1}{t} \left[\int_X |u| \phi^n + n \right] \leq \frac{|\sup_X u| + c}{t},
\end{aligned}$$

where we have used Proposition 1.7 and Corollary 2.3 from [GZ1] in the last two inequalities.

1.8. Proposition. *Let $u_j, u \in PSH(X, \emptyset)$. Then the following two statements are equivalent:*

- i) $u_j - u \rightarrow 0$ in C_X ;
- ii) $\sup_{j \geq 1} |\sup_X u_j| < +\infty$ and $\max(u_j, -t) - \max(u, -t) \rightarrow 0$ in C_X for every $t > 0$.

Proof. i) $\Rightarrow$ ii) By a result from [Hö] we have that $\int_X u_j \phi^n \rightarrow \int_X u \phi^n > -\infty$, $j \rightarrow \infty$.

Thus we obtain

$$\inf_{j \geq 1} \sup_X u_j \geq \inf_{j \geq 1} \int_X u_j \phi^n > -\infty.$$

Since $|u_j - u| \geq |\max(u_j, -t) - \max(u, -t)|$ we get

$$C_X(\{|\max(u_j, -t) - \max(u, -t)| > \delta\}) \leq C_X(\{|u_j - u| > \delta\}) \rightarrow 0,$$

for all $\delta > 0, t > 0$.

ii) $\Rightarrow$ i) We can assume that $u_j, u \leq 0$. Set

$$M = \sup_{j \geq 1} |\sup_X u_j| + |\sup_X u|$$

By Proposition 1.7 we have

$$\begin{aligned}
C_X(\{|u_j - u| > 3\delta\}) &\leq C_X(\{|u_j - \max(u_j, -t)| > \delta\}) + C_X(\{|\max(u, -t) - u| > \delta\}) \\
&+ C_X(\{|\max(u_j, -t) - \max(u, -t)| > \delta\}) \leq C_X(\{u_j < -t\}) + C_X(\{u < -t\}) \\
&+ C_X(\{|\max(u_j, -t) - \max(u, -t)| > \delta\}) \leq \frac{|\sup_X u_j| + |\sup_X u| + 2c}{t} \\
&+ C_X(\{|\max(u_j, -t) - \max(u, -t)| > \delta\}) \leq \frac{M + 2c}{t} \\
&+ C_X(\{|\max(u_j, -t) - \max(u, -t)| > \delta\})
\end{aligned}$$

for any $t > t_0 > 0, \delta > 0$. Hence

$$\overline{\lim}_{j \rightarrow \infty} C_X(\{|u_j - u| > 3\delta\}) \leq \frac{M + 2c}{t}.$$

Finally by letting $t \rightarrow +\infty$ we get

$$\lim_{j \rightarrow \infty} C_X(\{|u_j - u| > 3\delta\}) = 0$$

for every $\delta > 0$.

1.9. Proposition. *Let $u_j \in \mathcal{E}^-(X, \emptyset)$. Then the following two statements are equivalent:*

- i) $\inf_{j \geq 1} \sup_X u_j > -\infty$ and $\phi_{u_j}^n \ll C_X$ uniformly for $j \geq 1$;
- ii) $\lim_{t \rightarrow +\infty} \overline{\lim}_{j \rightarrow \infty} \int_{\{u_j \leq -t\}} \phi_{u_j}^n = 0$.

Proof. i) $\Rightarrow$ ii) This is a direct application of Proposition 1.7.

ii) $\Rightarrow$ i) We claim that $\inf_{j \geq 1} \sup_X u_j > -\infty$. Suppose on contrary that $\inf_{j \geq 1} \sup_X u_j = -\infty$.

Hence

$$\overline{\lim}_{j \rightarrow \infty} \int_{\{u_j \leq -t\}} \phi_{u_j}^n = \overline{\lim}_{j \rightarrow \infty} \int_X \phi_{u_j}^n = 1,$$

but this contradicts

$$\lim_{t \rightarrow +\infty} \overline{\lim}_{j \rightarrow \infty} \int_{\{u_j \leq -t\}} \phi_{u_j}^n = 0.$$

Let us fix $\epsilon > 0$. We choose t_0 and then $j_0 = j_0(t_0) > 1$ such that

$$\int_{\{u_j \leq -t_0\}} \phi_{u_j}^n < \epsilon,$$

for all $j \geq j_0$. For each Borel set $E \subset X$ we have

$$\begin{aligned} \int_E \phi_{u_j}^n &= \int_{E \cap \{u_j \leq -t_0\}} \phi_{u_j}^n + \int_{E \cap \{u_j > -t_0\}} \phi_{u_j}^n \leq \int_{\{u_j \leq -t_0\}} \phi_{u_j}^n + \int_{E \cap \{u_j > -t_0\}} \phi_{\max(u_j, -t_0)}^n \\ &\leq \int_{\{u_j \leq -t_0\}} \phi_{u_j}^n + t_0^n C_X(E) \leq \epsilon + t_0^n C_X(E) \end{aligned}$$

for all $j \geq j_0$. On the other hand, since $\sum_{k=1}^{j_0} \phi_{u_k}^n \ll C_X$ we can choose $\delta_1 > 0$ such that

$\sum_{k=1}^{j_0} \phi_{u_k}^n(E) < \epsilon$ for all Borel sets $E \subset X$ with $C_X(E) < \delta_1$. Hence $\phi_{u_j}^n(E) < 2\epsilon$ for all $j \geq 1$ and all Borel sets $E \subset X$, such that $C_X(E) < \delta = \min(\delta_1, \frac{\epsilon}{t_0^n})$.

The next proposition is a modified version of Lemma 2 in [Xi2]:

1.10. Proposition. *Let $u_j, v_j \in \mathcal{E}^-(X, \emptyset)$ be such that $u_j \geq v_j$ for $j \geq 1$. Assume that $\phi_{v_j}^n \ll C_X$ uniformly for $j \geq 1$. Let also $\inf_{j \geq 1} \sup_X v_j > -\infty$. Then $\phi_{u_j}^n \ll C_X$ uniformly for $j \geq 1$.*

Proof. By Theorem 1.5 in [GZ2] we have

$$\int_{\{u_j < -2t\}} \phi_{u_j}^n \leq 2^n \int_{\{v_j < \frac{u_j}{2} - t\}} \phi_{\frac{u_j}{2}}^n \leq 2^n \int_{\{v_j < \frac{u_j}{2} - t\}} \phi_{v_j}^n \leq 2^n \int_{\{v_j < -t\}} \phi_{v_j}^n,$$

for every $t > 0$. By Proposition 1.9 we obtain $\phi_{u_j}^n \ll C_X$ uniformly for $j \geq 1$.

We remark that the proof shows that the result still holds for any family of functions u_α and v_α , $\alpha \in \Lambda$ instead of just sequences.

1.11. Proposition. *Let $u_j \in \mathcal{E}(X, \emptyset)$, $u \in PSH(X, \emptyset)$ be such that $u_j \rightarrow u$ in C_X . Then the following two statements are equivalent:*

i) $u \in \mathcal{E}(X, \emptyset)$;

ii) $\phi_{u_j}^n \ll C_X$ uniformly for $j \geq 1$.

Proof. i) $\Rightarrow$ ii) By Proposition 1.6 we have

$$\begin{aligned} \int_{\{u_j \leq -t\}} \phi_{u_j}^n &= \int_{\{u_j \leq -t\}} \phi_{\max(u_j, -t)}^n \leq \int_{\{u \leq -t+1\}} \phi_{\max(u_j, -t)}^n + \int_{\{|u_j - u| > 1\}} \phi_{\max(u_j, -t)}^n \\ &\leq \int_{\{u \leq -t+1\}} \phi_{\max(u_j, -t)}^n + t^n(\{|u_j - u| > 1\}). \end{aligned}$$

By coupling the quasi-continuity of u (Theorem 3.5 in [BT2]) and the facts that $\phi_{\max(u_j, -t)}^n \ll C_X$ uniformly for $j \geq 1$ and $\phi_{\max(u, -t)}^n \ll C_X$ uniformly for $t > 0$ (see Proposition 1.10), we get

$$\overline{\lim}_{t \rightarrow +\infty} \overline{\lim}_{j \rightarrow \infty} \int_{\{u_j \leq -t\}} \phi_{u_j}^n \leq \overline{\lim}_{t \rightarrow +\infty} \int_{\{u \leq -t+1\}} \phi_{\max(u, -t)}^n = 0.$$

By Proposition 1.9 we obtain $\phi_{u_j}^n \ll C_X$ uniformly for $j \geq 1$.

ii) $\Rightarrow$ i) This is a part of Theorem 3 in [Xi2].

The last proposition is a somewhat stronger version of implication ii) $\Rightarrow$ i):

1.12. Proposition. *Let $u_j \in \mathcal{E}(X, \emptyset)$, $u \in PSH(X, \emptyset)$ be such that $u_j \rightarrow u$ in $L^1(X)$. Assume that $\phi_{u_j}^n \ll C_X$ uniformly for $j \geq 1$. Then $u \in \mathcal{E}(X, \emptyset)$.*

Proof. Set $v_j = \max(u_j, u) \in \mathcal{E}(X, \emptyset)$. By Hartogs' lemma (see [Hö]) we have $v_j \rightarrow u$ in C_X . Thus, by Proposition 1.10, we get that $\phi_{v_j}^n \ll C_X$ uniformly for $j \geq 1$. Now, using Proposition 1.11, we obtain $u \in \mathcal{E}(X, \emptyset)$.

2. CONVERGENCE IN CAPACITY Below we prove various stability results regarding convergence in capacity in the class $\mathcal{E}(X, \omega)$. We begin with a technical result which will be used later on. To authors' knowledge such a theorem is new although some similar ideas can be found in Lemma 2.3 in [H1] and in Theorem 2 in [Xi2].

2.1. Theorem. *Let $u_j, v_j \in \mathcal{E}(X, \emptyset)$ be such that*

i) $\phi_{u_j}^n \ll C_X$ uniformly for $j \geq 1$;

ii) $\lim_{j \rightarrow \infty} \int_{\{u_j < v_j - \delta\}} \phi_{u_j}^n = 0$ for all $\delta > 0$.

Then $\lim_{j \rightarrow \infty} C_X(\{u_j < v_j - \delta\}) = 0$ for all $\delta > 0$.

Proof. We can assume, by adding constants to both u_j, v_j if necessary, that $\sup_X u_j = 0$ for all $j \geq 1$. Set $u_{jt} := \max(u_j, -t)$. For each $k = 0, \dots, n$ we will prove inductively that

$$(1) \quad \lim_{j \rightarrow \infty} \sup \left\{ \int_{\{u_j < v_j - \delta\}} \phi_{u_j}^{n-k} \wedge \phi_{\varphi}^k : \varphi \in PSH(X, \emptyset), -1 \leq \varphi \leq 0 \right\} = 0$$

for every $\delta > 0$. If $k = 0$ then (1) holds by assumption.

Assume that (1) holds for $k - 1$. We will prove that

$$\lim_{j \rightarrow \infty} \sup \left\{ \int_{\{u_j < v_j - 3\delta\}} \phi_{u_j}^{n-k} \wedge \phi_{\varphi}^k : \varphi \in PSH(X, \emptyset), -1 \leq \varphi \leq 0 \right\} = 0$$

for any $\delta > 0$. We fix $t \geq 1$ and $\varphi \in PSH(X, \emptyset)$ satisfying $-1 \leq \varphi \leq 0$. Then, by Theorem 2.3 in [Di3], Corollary 1.7 in [GZ2] and Proposition 1.6 we have

$$\begin{aligned}
& \int_{\{u_j < v_j - 3\delta\}} \phi_{u_j}^{n-k} \wedge \phi_\varphi^k \leq \int_{\{u_j + \frac{\delta}{t} u_{jt} < v_j + \frac{\delta}{t} \varphi - 2\delta\}} \phi_{u_j}^{n-k} \wedge \phi_\varphi^k \\
& \leq \frac{t + \delta}{\delta} \int_{\{u_j + \frac{\delta}{t} u_{jt} < v_j + \frac{\delta}{t} \varphi - 2\delta\}} \phi_{\frac{v_j + \frac{\delta}{t} \varphi - 2\delta}{1 + \frac{\delta}{t}}} \wedge \phi_{u_j}^{n-k} \wedge \phi_\varphi^{k-1} \\
& \leq \frac{t + \delta}{\delta} \int_{\{u_j + \frac{\delta}{t} u_{jt} < v_j + \frac{\delta}{t} \varphi - 2\delta\}} \phi_{\frac{u_j + \frac{\delta}{t} u_{jt}}{1 + \frac{\delta}{t}}} \wedge \phi_{u_j}^{n-k} \wedge \phi_\varphi^{k-1} \leq \frac{t}{\delta} \int_{\{u_j < v_j - \delta\}} \phi_{u_j}^{n-k+1} \wedge \phi_\varphi^{k-1} \\
& + \int_{\{u_j < v_j - \delta\}} \phi_{u_{jt}} \wedge \phi_{u_j}^{n-k} \wedge \phi_\varphi^{k-1} \leq \frac{t + \delta}{\delta} \int_{\{u_j < v_j - \delta\}} \phi_{u_j}^{n-k+1} \wedge \phi_\varphi^{k-1} \\
& + \int_{\{u_j \leq -t\}} \phi_{u_{jt}} \wedge \phi_{u_j}^{n-k} \wedge \phi_\varphi^{k-1} \leq \frac{t + \delta}{\delta} \int_{\{u_j < v_j - \delta\}} \phi_{u_j}^{n-k+1} \wedge \phi_\varphi^{k-1} + \int_{\{u_j \leq -t\}} \phi_{u_j}^{n-k+1} \wedge \phi_\varphi^{k-1}.
\end{aligned}$$

By the induction hypothesis we get

$$\begin{aligned}
& \overline{\lim}_{j \rightarrow \infty} \sup \left\{ \int_{\{u_j < v_j - 3\delta\}} \phi_{u_j}^{n-k} \wedge \phi_\varphi^k : \varphi \in PSH(X, \emptyset), -1 \leq \varphi \leq 0 \right\} \\
& \leq \sup \left\{ \int_{\{u_j \leq -t\}} \phi_{u_j}^{n-k+1} \wedge \phi_\varphi^{k-1} : \varphi \in PSH(X, \emptyset), -1 \leq \varphi \leq 0, j \geq 1 \right\},
\end{aligned}$$

for every $t \geq 1$.

Note that $\frac{u_j + \varphi}{2} \geq u_j - 1$, thus by Proposition 1.10 we get that $\phi_{\frac{u_j + \varphi}{2}}^n \ll C_X$ uniformly for $j \geq 1$. Expanding the left hand side we obtain

$$\phi_{u_j}^{n-k+1} \wedge \phi_\varphi^{k-1} \ll C_X \text{ uniformly for } j \geq 1.$$

Now by letting $t \rightarrow +\infty$ (and using $\phi_{u_j}^{n-k+1} \wedge \phi_\varphi^{k-1} \ll C_X$ uniformly for $j \geq 1$), from Proposition 1.7 we obtain

$$\lim_{j \rightarrow \infty} \sup \left\{ \int_{\{u_j < v_j - 3\delta\}} \phi_{u_j}^{n-k} \wedge \phi_\varphi^k : \varphi \in PSH(X, \emptyset), -1 \leq \varphi \leq 0 \right\} = 0.$$

Theorem 2.1 has various consequences regarding stability of the Monge-Ampère operator. Below we list some of them.

As a direct corollary we obtain the following theorem:

2.2. Theorem. *Let $u_j, v_j \in \mathcal{E}(X, \emptyset)$ be such that*

- i) $\phi_{u_j}^n + \phi_{v_j}^n \ll C_X$ uniformly for $j \geq 1$;
- ii) $\lim_{j \rightarrow \infty} \left[\int_{\{u_j < v_j - \delta\}} \phi_{u_j}^n + \int_{\{v_j < u_j - \delta\}} \phi_{v_j}^n \right] = 0$ for all $\delta > 0$.

Then $u_j - v_j \rightarrow 0$ in C_X .

Recently the second named author obtained characterization of convergence in capacity for bounded $\emptyset$ -psh functions (Theorem 2.1 in [H1]). Our next result provides such a characterization in the class $\mathcal{E}(X, \emptyset)$:

2.3. Theorem. *Let $u_j \in \mathcal{E}(X, \emptyset)$ and $u \in PSH(X, \emptyset)$. Then the following three statements are equivalent:*

- i) $u \in \mathcal{E}(X, \emptyset)$ and $u_j \rightarrow u$ in C_X ;
- ii) $\phi_{u_j}^n \ll C_X$ uniformly for $j \geq 1$ and $u_j \rightarrow u$ in C_X ;

iii) $\phi_{u_j}^n \ll C_X$ uniformly for $j \geq 1$, $\overline{\lim}_{j \rightarrow \infty} u_j \leq u$ and $\lim_{j \rightarrow \infty} \int_{\{u_j < u - \delta\}} \phi_{u_j}^n = 0$ for all $\delta > 0$.

Proof. i) $\Leftrightarrow$ ii) This equivalence is the content of Proposition 1.11.

ii) $\Rightarrow$ iii) This implication is trivial.

iii) $\Rightarrow$ i) This is a direct application Theorem 2.1, coupled with Proposition 1.11.

Recall that the weak convergence of Monge-Ampère measures does not imply the weak convergence of the corresponding functions (and vice versa). Thus such convergence problems are considered with additional restrictions. In [CK2] good convergence properties were obtained (in the setting of domains in $\mathbf{C}^n$) under the assumption that all the Monge-Ampère measures are dominated by a fixed measure vanishing on pluripolar sets. In the Kähler setting the corresponding problem was studied by Xing in [Xi2]. He worked in a subclass $\mathcal{E}^1(X, \phi) \subset \mathcal{E}(X, \phi)$ however all his arguments can be applied in the whole $\mathcal{E}(X, \phi)$ provided one has an uniqueness (modulo additive constant) for the solutions of the Monge-Ampère equation

$$\phi_\varphi = \mu, \quad \varphi \in \mathcal{E}(X, \phi),$$

where μ is any positive Borel measure satisfying $\int_X d\mu = \int_X \phi^n$ and vanishing on pluripolar sets. This uniqueness statement was recently obtained in [Di3]. Thus coupling Xing's arguments together with Theorem 2.2 one gets the following stability theorem:

2.4. Theorem. *Let $u_j \in \mathcal{E}(X, \phi)$ and $u \in \text{PSH}(X, \phi)$. Assume that $\omega_{u_j}^n \leq d\mu$ for some measure $d\mu \ll C_X$ (equivalently: μ does not charge pluripolar sets). Let also $\sup_X u_j = \sup_X u$. Then the following three statements are equivalent:*

- i) $u_j \rightarrow u$ in C_X .
- ii) $u_j \rightarrow u$ in L^1 .
- iii) $\omega_{u_j}^n \rightarrow \omega_u^n$ weakly.

The implication ii) $\Leftrightarrow$ iii) is essentially due to Xing. In particular Corollary 1 in [Xi2] (and Proposition 1.11) yields ii) $\Rightarrow$ iii). For the other implication one can proceed exactly as in Theorem 8 in [Xi2]: one can extract a subsequence from u_j convergent in L^1 to some function v . It follows that $\phi_u^n = \phi_v^n$, so by uniqueness (now in $\mathcal{E}(X, \phi)$) $u = v$. Thus, since any subsequence has subsequence convergent in L^1 to u one concludes that $u_j \rightarrow u$ in L^1 .

Since implication i) $\Rightarrow$ ii) is trivial, we only have to show that ii) implies i). But we know that $\phi_{u_j}^n \ll C_X$ (since all the measures are dominated by μ). By [Ce3] (see also Lemma 1.4 in [CK2]) we know that

$$\forall M \in \mathbf{R} \quad \lim_{j \rightarrow \infty} \int_X \max(u_j, M) - \max(u, M) d\mu = 0.$$

Thus we obtain

$$\int_{\{u_j < u - \delta\}} \phi_{u_j}^n + \int_{\{u < u_j - \delta\}} \phi_u^n \leq \int_{\{u_j < M\} \cup \{u < M\}} d\mu + \frac{1}{\delta} \int_X |\max(u_j, M) - \max(u, M)| d\mu.$$

If M is sufficiently negative by the assumption $\mu \ll C_X$ and proposition 1.7 we obtain that the first term on the right hand side can be made arbitrarily small (independently of j). Finally we obtain

$$\lim_{j \rightarrow \infty} \left[\int_{\{u_j < u - \delta\}} \phi_{u_j}^n + \int_{\{u < u_j - \delta\}} \phi_u^n \right] = 0.$$

Thus by theorem 2.2 $u_j \rightarrow u$ in C_X .

Our next result concerns a new type of stability. Instead of controlling the deviation of u_j from v_j we impose an assumption on the variation of their Monge-Ampère measures. Such a theorem is new even for bounded ϕ -psh functions.

2.5. Theorem. *Let $u_j, v_j \in \mathcal{E}(X, \phi)$ be such that*

- i) $\lim_{j \rightarrow \infty} |\sup_X u_j - \sup_X v_j| = 0$;
- ii) $\phi_{u_j}^n + \phi_{v_j}^n \ll C_X$ uniformly for $j \geq 1$;
- iii) $\lim_{j \rightarrow \infty} \int_X |\phi_{u_j}^n - \phi_{v_j}^n| = 0$.

Then $u_j - v_j \rightarrow 0$ in C_X .

Before we start the proof we state an auxiliary lemma. To authors' knowledge such an estimate is new and might be of independent interest.

2.6. Lemma. *Let $u, v \in \mathcal{E}(X, \phi)$. Then*

$$\int_X |\phi_u^k \wedge \phi_v^{n-k} - \phi_u^n| \leq 2 \left[\int_X |\phi_u^n - \phi_v^n| \right]^{\frac{n}{n-k}},$$

Proof. Set

$$\mu = \frac{1}{2}[\phi_u^n + \phi_v^n].$$

We choose $f, g \in L^1(d\mu)$, $f, g \geq 0$ such that

$$\phi_u^n = f d\mu, \quad \phi_v^n = g d\mu.$$

By Theorem 2.1 in [Di3] and Hölder inequality we get

$$\begin{aligned} \int_E \phi_u^k \wedge \phi_v^{n-k} - \int_E \phi_u^n &\geq \int_E f^{\frac{k}{n}} g^{\frac{n-k}{n}} d\mu - \int_E f d\mu = \int_E f^{\frac{k}{n}} [g^{\frac{n-k}{n}} - f^{\frac{n-k}{n}}] d\mu \\ &\geq - \int_E f^{\frac{k}{n}} |g^{\frac{n-k}{n}} - f^{\frac{n-k}{n}}| d\mu \geq - \int_E f^{\frac{k}{n}} |g - f|^{\frac{n-k}{n}} d\mu \geq - \left[\int_E f d\mu \right]^{\frac{k}{n}} \left[\int_E |g - f| d\mu \right]^{\frac{n-k}{n}} \\ &\geq - \left[\int_E |g - f| d\mu \right]^{\frac{n-k}{n}} \geq - \left[\int_X |\phi_u^n - \phi_v^n| \right]^{\frac{n-k}{n}} \end{aligned}$$

for any Borel set $E \subset X$. Similarly we get

$$\int_{X \setminus E} \phi_u^k \wedge \phi_v^{n-k} - \int_{X \setminus E} \phi_u^n \geq - \left[\int_X |\phi_u^n - \phi_v^n| \right]^{\frac{n-k}{n}}$$

for any Borel set $E \subset X$. Hence

$$\int_E \phi_u^k \wedge \phi_v^{n-k} - \int_E \phi_u^n \leq \left[\int_X |\phi_u^n - \phi_v^n| \right]^{\frac{n-k}{n}}$$

for all Borel set $E \subset X$. Combination of these inequalities yields

$$\left| \int_E \phi_u^k \wedge \phi_v^{n-k} - \int_E \phi_u^n \right| \leq \left[\int_X |\phi_u^n - \phi_v^n| \right]^{\frac{n-k}{n}},$$

for all Borel $E \subset X$. This implies that

$$\int_X |\phi_u^k \wedge \phi_v^{n-k} - \phi_u^n| \leq 2 \left[\int_X |\phi_u^n - \phi_v^n| \right]^{\frac{n}{n-k}}.$$

Proof of Theorem 2.5.

Case I: $u_j, v_j \in PSH(X, \omega) \cap L^\infty(X)$. Set

$$a_j = [\sup\{\int_X ||\phi_{u_j}^k \wedge \phi_{v_j}^{n-k} - \phi_{u_j}^m \wedge \phi_{v_j}^{n-m}|| : 1 \leq k, m \leq n\}]^{\frac{1}{4}}.$$

By Lemma 2.6 we get $\lim_{j \rightarrow \infty} a_j = 0$. We shall prove that

$$(2) \quad \lim_{j \rightarrow \infty} \sup_{t \in \mathbf{R}} \left\{ \int_{\{|u_j - v_j - t| \leq a_j\}} \phi_{u_j}^n \right\} = 1.$$

Suppose that there exists $\epsilon_0 > 0$ such that

$$\int_{\{|u_j - v_j - t| \leq a_j\}} \phi_{u_j}^n \leq 1 - 2\epsilon_0$$

for all $t \in \mathbf{R}$, $j \geq 1$. Set

$$t_j = \sup\{t \in \mathbf{R} : \int_{\{u_j < v_j + t + a_j\}} \phi_{u_j}^n \leq 1 - \epsilon_0\}$$

Replacing $v_j + t_j$ by v_j we can assume that $t_j = 0$. Then $\int_{\{u_j < v_j + a_j\}} \phi_{u_j}^n \leq 1 - \epsilon_0$ and

$\int_{\{u_j \leq v_j + a_j\}} \phi_{u_j}^n \geq 1 - \epsilon_0$. Hence

$$\begin{aligned} \int_{\{v_j < u_j + a_j\}} \phi_{u_j}^n &= 1 - \int_{\{u_j + a_j \leq v_j\}} \phi_{u_j}^n = 1 - \int_{\{u_j \leq v_j + a_j\}} \phi_{u_j}^n \\ &+ \int_{\{v_j - a_j < u_j \leq v_j + a_j\}} \phi_{u_j}^n \leq 1 - \epsilon_0. \end{aligned}$$

Since $\int_{\{|u_j - v_j| \leq a_j\}} \phi_{u_j}^n \leq 1$ we can choose $s_j \in [-a_j + a_j^2, a_j - a_j^2]$ satisfying

$$\int_{\{|u_j - v_j - s_j| < a_j^2\}} \phi_{u_j}^n \leq 2a_j.$$

Replacing $v_j + s_j$ by v_j we can assume that $s_j = 0$. One easily obtains the following inequalities

$$\int_{\{u_j < v_j + a_j^2\}} \phi_{u_j}^n \leq 1 - \epsilon_0, \quad \int_{\{v_j < u_j + a_j^2\}} \phi_{u_j}^n \leq 1 - \epsilon_0, \quad \int_{\{|u_j - v_j| < a_j^2\}} \phi_{u_j}^n \leq 2a_j.$$

By [GZ2] we can find $\rho_j \in \mathcal{E}(X, \omega)$, such that $\sup_X \rho_j = 0$ and $\omega_{\rho_j}^n = \frac{1}{1-\epsilon_0} 1_{\{u_j < v_j\}} \omega_{u_j}^n + c_j 1_{\{u_j \geq v_j\}} \omega_{u_j}^n$ ($c_j \geq 0$ is chosen so that the measure has total mass 1). Set

$$U_j = \{(1 - a_j^3)u_j < (1 - a_j^3)v_j + a_j^3 \rho_j\} \subset \{u_j < v_j\}.$$

By Theorem 2.1 in [Di3] we get

$$\omega_{u_j}^{n-1} \wedge \omega_{(1-a_j^3)v_j + a_j^3 \rho_j} \geq (1 - a_j^3) \omega_{u_j}^{n-1} \wedge \omega_{v_j} + \frac{a_j^3}{(1 - \epsilon_0)^{\frac{1}{n}}} \omega_{u_j}^n$$

on U_j . By Theorem 2.3 in [Di3] we obtain

$$\begin{aligned}
& (1 - a_j^3) \int_{U_j} \omega_{u_j}^{n-1} \wedge \omega_{v_j} + \frac{a_j^3}{(1 - \epsilon_0)^{\frac{1}{n}}} \int_{U_j} \omega_{u_j}^n \leq \int_{U_j} \omega_{(1-a_j^3)v_j + a_j^3 \rho_j} \wedge \omega_{u_j}^{n-1} \\
& \leq \int_{U_j} \omega_{(1-a_j^3)u_j} \wedge \omega_{u_j}^{n-1} = (1 - a_j^3) \int_{U_j} \omega_{u_j}^n + a_j^3 \int_{U_j} \omega \wedge \omega_{u_j}^{n-1} \\
& \leq (1 - a_j^3) \left(\int_{U_j} \omega_{u_j}^{n-1} \wedge \omega_{v_j} + a_j^4 \right) + a_j^3 \int_{U_j} \omega \wedge \omega_{u_j}^{n-1}.
\end{aligned}$$

Hence

$$\begin{aligned}
& \frac{1}{(1 - \epsilon_0)^{\frac{1}{n}}} \left[\int_{\{u_j \leq v_j - a_j^2\}} \omega_{u_j}^n - \int_{\{\rho_j \leq -\frac{1-a_j^3}{a_j}\}} \omega_{u_j}^n \right] \leq \frac{1}{(1 - \epsilon_0)^{\frac{1}{n}}} \int_{U_j} \omega_{u_j}^n \\
& \leq a_j + \int_{U_j} \omega \wedge \omega_{u_j}^{n-1} \leq a_j + \int_{\{u_j < v_j\}} \omega \wedge \omega_{u_j}^{n-1}.
\end{aligned}$$

Similarly to ρ_j we define $\vartheta_j \in \mathcal{E}(X, \emptyset)$, such that $\sup_X \vartheta_j = 0$ and $\omega_{\vartheta_j}^n = \frac{1}{1-\epsilon_0} 1_{\{v_j < u_j\}} \omega_{u_j}^n + d_j 1_{\{v_j \geq u_j\}} \omega_{u_j}^n$ (d_j plays the same role as c_j above). Set

$$V_j = \{(1 - a_j^3)v_j < (1 - a_j^3)u_j + a_j^3 \vartheta_j\} \subset \{v_j < u_j\}.$$

Similarly we get

$$\begin{aligned}
& \frac{1}{(1 - \epsilon_0)^{\frac{1}{n}}} \left[\int_{\{v_j \leq u_j - a_j^2\}} \omega_{u_j}^n - \int_{\{\vartheta_j \leq -\frac{1-a_j^3}{a_j}\}} \omega_{u_j}^n \right] \leq \frac{1}{(1 - \epsilon_0)^{\frac{1}{n}}} \int_{V_j} \omega_{u_j}^n \\
& \leq a_j + \int_{V_j} \omega \wedge \omega_{u_j}^{n-1} \leq a_j + \int_{\{v_j < u_j\}} \omega \wedge \omega_{u_j}^{n-1}.
\end{aligned}$$

Combination of these inequalities yields

$$\begin{aligned}
& \frac{1}{(1 - \epsilon_0)^{\frac{1}{n}}} [1 - 2a_j - 2 \int_{\{\rho_j \leq -\frac{1-a_j^3}{a_j}\}} \omega_{u_j}^n] \leq \frac{1}{(1 - \epsilon_0)^{\frac{1}{n}}} \left[\int_{\{|u_j - v_j| \geq a_j^2\}} \omega_{u_j}^n - 2 \int_{\{\rho_j \leq -\frac{1-a_j^3}{a_j}\}} \omega_{u_j}^n \right] \\
& \leq 2a_j + 1.
\end{aligned}$$

Now, by Proposition 1.7 and assumption ii), if we let $j \rightarrow \infty$ we would obtain

$$\frac{1}{(1 - \epsilon_0)^{\frac{1}{n}}} \leq 1,$$

a contradiction.

Using (2) we can choose $k_j \in \mathbf{R}$ such that

$$\lim_{j \rightarrow \infty} \int_{\{|u_j - v_j - k_j| > a_j\}} \phi_{u_j}^n = 0.$$

Moreover from iii) we get

$$\lim_{j \rightarrow \infty} \int_{\{|u_j - v_j - k_j| > a_j\}} \phi_{v_j}^n = 0.$$

By Theorem 2.2 we obtain $u_j - v_j - k_j \rightarrow 0$ in C_X . Moreover from i) we get $\lim_{j \rightarrow \infty} k_j = 0$. Hence $u_j - v_j \rightarrow 0$ in C_X .

Case II: $u_j, v_j \in \mathcal{E}(X, \omega)$. We can assume, adding constants if necessary, that $\sup_X u_j = \sup_X v_j = 0$. Choose $t_j \rightarrow +\infty$ such that $\lim_{j \rightarrow \infty} \int_{\{u_j \leq -t_j\}} \omega_{u_j}^n + \int_{\{v_j \leq -t_j\}} \omega_{v_j}^n = 0$. Using Case I for $\max(u_j, -t_j)$ and $\max(v_j, -t_j)$ we obtain $\max(u_j, -t_j) - \max(v_j, -t_j) \rightarrow 0$ in C_X . Therefore $u_j - v_j \rightarrow 0$ in C_X .

Our last result is a generalization of Theorem 5 from [Xi2] and Theorem 3.4 from [Ko2]. Kołodziej proved the statement for bounded functions, while Xing needed the additional assumption that $\forall j \geq 1, v_j \geq v_0$ for some fixed function $v_0 \in \mathcal{E}(X, \phi)$. In fact we show that this condition is superfluous.

2.7. Theorem. *Let $u_j, v_j, v \in \mathcal{E}(X, \phi)$, $u \in PSH(X, \phi)$ and $A > 1$ be such that $u_j \rightarrow u$ in $L^1(X)$ and $\phi_{u_j}^n \leq A\phi_{v_j}^n$ for $j \geq 1$. Assume that $v_j \rightarrow v$ in C_X . Then $u \in \mathcal{E}(X, \phi)$ and $u_j \rightarrow u$ in C_X .*

Proof. For each $t > 0$ we set

$$u_{jt} := \max(u_j, -t), \quad u_t := \max(u, -t), \quad v_{jt} := \max(v_j, -t), \quad v_t := \max(v, -t),$$

$$T_{jt} := \sum_{k=0}^{n-1} \phi_{v_{jt}}^k \wedge \phi_{v_t}^{n-1-k}.$$

We have

$$\int_{\{u_j < u - \delta\}} \phi_{u_j}^n \leq A \int_{\{u_j < u - \delta\}} \phi_{v_j}^n.$$

The latter integral can be estimated by

$$\begin{aligned} \int_{\{u_j < u - \delta\}} \phi_{v_j}^n &\leq \int_{\{u_j \leq -t\} \cup \{u \leq -t\} \cup \{v_j \leq -t\}} \phi_{v_j}^n + \int_{\{u_{jt} < u_t - \delta\}} \phi_{v_{jt}}^n \leq \int_{\{u_j \leq -t\} \cup \{u \leq -t\} \cup \{v_j \leq -t\}} \phi_{v_j}^n \\ &+ \frac{1}{\delta} \int_X |u_t - u_{jt}| \phi_{v_{jt}}^n \leq \int_{\{u_j \leq -t\} \cup \{u \leq -t\} \cup \{v_j \leq -t\}} \phi_{v_j}^n + \frac{1}{\delta} \int_X |u_t + \epsilon - u_{jt}| \phi_{v_{jt}}^n \\ &+ \frac{\epsilon}{\delta} \leq \int_{\{u_j \leq -t\} \cup \{u \leq -t\} \cup \{v_j \leq -t\}} \phi_{v_j}^n + \frac{1}{\delta} \int_X (u_t - u_{jt}) \phi_{v_{jt}}^n + \frac{t + \epsilon}{\delta} \int_{\{u_{jt} > u_t + \epsilon\}} \phi_{v_{jt}}^n + \frac{2\epsilon}{\delta} \\ &\leq \int_{\{u_j \leq -t\} \cup \{u \leq -t\} \cup \{v_j \leq -t\}} \phi_{v_j}^n + \frac{1}{\delta} \int_X (u_t - u_{jt}) (\phi_{v_{jt}}^n - \phi_{v_t}^n) + \frac{1}{\delta} \int_X (u_t - u_{jt}) \phi_{v_t}^n \end{aligned}$$

$$\begin{aligned}
& + \frac{t+\epsilon}{\delta} \int_{\{u_{jt} > u_t + \epsilon\}} \phi_{v_{jt}}^n + \frac{2\epsilon}{\delta} = \int_{\{u_j \leq -t\} \cup \{u \leq -t\} \cup \{v_j \leq -t\}} \phi_{v_j}^n + \frac{1}{\delta} \int_X (v_t - v_{jt})(\phi_{u_{jt}} - \phi_{u_t}) \wedge T_{jt} \\
& + \frac{1}{\delta} \int_X (u_t - u_{jt})\phi_{v_t}^n + \frac{t+\epsilon}{\delta} \int_{\{u_{jt} > u_t + \epsilon\}} \phi_{v_{jt}}^n + \frac{2\epsilon}{\delta} \leq \int_{\{u_j \leq -t\} \cup \{u \leq -t\} \cup \{v_j \leq -t\}} \phi_{v_j}^n \\
& + \frac{1}{\delta} \int_X |v_t - v_{jt}|(\phi_{u_{jt}} + \phi_{u_t}) \wedge T_{jt} + \frac{1}{\delta} \int_X (u_t - u_{jt})\phi_{v_t}^n + \frac{t+\epsilon}{\delta} \int_{\{u_{jt} > u_t + \epsilon\}} \phi_{v_{jt}}^n + \frac{2\epsilon}{\delta}
\end{aligned}$$

for all $t > 0$ and all $\epsilon > 0$.

Observe that the second term goes to 0 as $j \rightarrow \infty$, since the integrated function converges to 0 in capacity and all the measures are uniformly absolutely continuous with respect to C_X for $j \geq 1$ (and fixed t). By [Ce3] we also get

$$\lim_{j \rightarrow \infty} \int_X (u_t - u_{jt})\phi_{v_t}^n = 0.$$

Note that Hartogs' lemma (see [Hö]) yields $C_X(\{u_{jt} > u_t + \epsilon\}) \rightarrow 0$ as $j \rightarrow \infty$. Moreover, since $\omega_{v_{jt}}^n \ll C_X$ uniformly for $j \geq 1$ we obtain

$$\lim_{j \rightarrow \infty} \int_{\{u_{jt} > u_t + \epsilon\}} \phi_{v_{jt}}^n = 0.$$

Coupling all these and letting $j \rightarrow \infty$ we get

$$\overline{\lim}_{j \rightarrow \infty} \int_{\{u_j < u - \delta\}} \phi_{u_j}^n \leq \sup_{j \geq 1} \int_{\{u_j \leq -t\} \cup \{u \leq -t\} \cup \{v_j \leq -t\}} \phi_{v_j}^n + \frac{2\epsilon}{\delta},$$

for all $t > 0$, $\epsilon > 0$. Letting $\epsilon \rightarrow 0$ and $t \rightarrow +\infty$ we obtain

$$\lim_{j \rightarrow \infty} \int_{\{u_j < u - \delta\}} \phi_{u_j}^n = 0.$$

By Theorem 2.3 we obtain that $u_j \rightarrow u$ in C_X .

REFERENCES

- [Bl] Z. Błocki, *Uniqueness and stability for the complex Monge-Ampère equation on compact Kähler manifolds*, Indiana Univ. Math. J. **52** (2003), no. 6, 1697-1701.
- [BT1] E. Bedford and B. A. Taylor, *The Dirichlet problem for the complex Monge-Ampère operator*. Invent. Math. **37** (1976), 1-44.
- [BT2] E. Bedford and B. A. Taylor, *A new capacity for plurisubharmonic functions*. Acta Math. **149** (1982), 1-40.
- [Ce1] U. Cegrell, *Pluricomplex energy*, Acta Math. **180** (1998), 187-217.
- [Ce2] U. Cegrell, *The general definition of the complex Monge-Ampère operator*, Ann. Inst. Fourier (Grenoble) **54** (2004), 159-179.
- [Ce3] U. Cegrell, *Convergence in capacity*, Technical report, Isaac Newton Institute for Mathematical Sciences, 2001.
- [CK1] U. Cegrell, S. Kolodziej, *The Dirichlet problem for the complex Monge-Ampère operator: Perron classes and rotation invariant measures*, Mich. Math. J. **41** (1994), 563-569.

- [CK2] U. Cegrell, S. Kołodziej, *The equation of complex Monge-Ampère type and stability of solutions*, Math. Ann. **334** (2006), 713-729.
- [CGZ] D. Coman, V. Guedj and A. Zeriahi, *Domains of definition of Monge-Ampère operators on compact Kähler manifolds*, Math. Zeit. **259** (2008), 393-418.
- [Di1] S. Dinew, *Cegrell classes on compact Kähler manifolds*, Ann. Polon. Math. **91** (2007), 179-195.
- [Di2] S. Dinew, *An inequality for mixed Monge-Ampère measures*, Math. Zeit. **262** (2009), 1-15.
- [Di3] S. Dinew, *Uniqueness in $\mathcal{E}(X, \phi)$* , J. Funct. Anal. **256** (2009), 2113-2122.
- [GZ1] V. Guedj and A. Zeriahi, *Intrinsic capacities on compact Kähler manifolds*, J. Geom. Anal. **15** (2005), no. 4, 607-639.
- [GZ2] V. Guedj and A. Zeriahi, *The weighted Monge-Ampère energy of quasiplurisubharmonic functions*, J. Funct. Anal. **250** (2007), 442-482.
- [H1] P. Hiep, *On the convergence in capacity on compact Kähler manifolds and its applications*, Proc. Amer. Math. Soc. **136** (2008), 2007-2018.
- [H2] P. Hiep, *Convergence in capacity*, Ann. Polon. Math. **93** (2008), 91-99.
- [Hö] L. Hörmander, *Notions of Convexity*, Progress in Mathematics **127**, Birkhäuser, Boston, (1994).
- [KH] N. Khue and P. Hiep, *A comparison principle for the complex Monge-Ampère operator in Cegrell's classes and applications*, to appear in Trans. Amer. Math. Soc.
- [Ko1] S. Kołodziej, *The Monge-Ampère equation on compact Kähler manifolds*, Indiana Univ. Math. J. **52** (2003), 667-686.
- [Ko2] S. Kołodziej, *The set of measures given by bounded solutions of the complex Monge-Ampère equation on compact Kähler manifolds*, J. London Math. Soc. **72** (2) (2005), 225-238.
- [Xi1] Y. Xing, *Continuity of the complex Monge-Ampère operator*, Proc. Amer. Math. Soc. **124** (1996), 457-467.
- [Xi2] Y. Xing, *Continuity of the Complex Monge-Ampère Operator on Compact Kähler Manifolds*, to appear in Math. Zeit. (<http://www.arxiv.org/>).

Jagiellonian University,
 Mathematics Institute,
 30-348 Łojasiewicza 6,
 Kraków, Poland
 slawomir.dinew@im.uj.edu.pl

Department of Mathematics
 University of Education (Dai hoc Su Pham Ha Noi)
 Cau Giay, Hanoi, Vietnam
 phhiep_vn@yahoo.com